## STANDARDIZED VENOM TESTING DATA



PATIENT NAME:	
DATE OF TESTING:	CHART#:
TREATING PHYSICIAN:	

Control Test Record	Histamine Control	Negative Control
WHEAL		
FLARE		

When mixing venom for testing purposes please make sure you rock back and forth OR swirl to mix; NEVER shake to mix. ALWAYS reconstitute and dilute venom with Sterile Albumin Saline with Phenol (0.4%). For complete information on these products, and their proper usage consult the package insert; available in the product packaging.







Standardized Venoms	Prick <b>1</b> μg,		Intradermal <b>0.0001</b> μg/mL		Intradermal <b>0.001</b> μg/mL		Intradermal <b>0.01</b> µg/mL		Intradermal <b>0.1</b> µg/mL		<b>Intradermal</b> <b>1</b> μg/mL	
RECORD BOTH WHEAL AND FLARE	W	F	W	F	W	F	W	F	W	F	W	F
Honey Bee Venom						:				•	:	
White-Faced Hornet Venom Protein						<u>:</u>				•	•	
Yellow Hornet Venom Protein						<u>:</u>				•	:	
Wasp Venom Protein						: :				•	:	
Yellow Jacket Venom Protein						:				•	:	

REFER TO PACKAGE INSERT FOR RECOMMENDED DOSAGE AND ADMINISTRATION

hsallergy.com