



## ALLERGENIC EXTRACTS

## Mite Extracts

Most homes in the United States have detectable levels of dust mite allergen in at least one bed.<sup>1</sup> Although there are ways to help limit exposure to dust mites through prevention techniques, immunotherapy treatment remains one of the safest ways to prevent an allergic reaction.

Ensure that your clinic is ready to diagnose and treat patients who suffer from dust mite allergies.

At HollisterStier Allergy, we grow, harvest, and manufacture our Standardized Dust Mite Extract at our Spokane, WA facility.

**Customizable Treatment for Your Practice Needs**

- Our standardized mite extracts are available in two distinct strengths, 10,000 AU/mL and 30,000 AU/mL, allowing you to customize the treatment to the patient for personalized care.
- Our higher concentration extract, 30,000 AU/mL, helps ensure an accurate diagnosis during skin testing for patients who might have been missed with conventional methods.<sup>2</sup>

ITEM NO.	DESCRIPTION	UNIT
6691	Standardized 50/50 Mite Mix, <i>D. pteronyssinus</i> and <i>D. farina</i>	5,000 AU/mL or 15,000 AU/mL (each species)
6692	Standardized Mite, <i>D. pteronyssinus</i>	10,000 AU/mL or 30,000 AU/mL
6720	Standardized Mite, <i>D. farina</i>	10,000 AU/mL or 30,000 AU/mL

**WARNING: ANAPHYLAXIS**

Allergenic extracts can cause anaphylaxis, including anaphylactic shock and death.

Emergency measures and healthcare providers trained in their use must be available immediately in the event of a life-threatening reaction. Observe individuals for at least 30 minutes following administration.

Do not administer to individuals with severe, unstable or uncontrolled asthma, history of severe systemic reaction to the allergen extract, or with medical conditions that reduce the ability to survive anaphylaxis.

Individuals with extreme sensitivity to these products, on an accelerated immunotherapy build-up, switching to another lot, receiving high doses of these products, or exposed to excessive amounts of the allergen may be at increased risk of anaphylaxis.

These products may not be suitable for individuals who may be unresponsive to epinephrine or inhaled bronchodilators, such as those taking beta-blockers.

Adverse reactions on our products can be reported by calling 800-495-7437, or by emailing [adversereactions@jubi.com](mailto:adversereactions@jubi.com).

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 800-FDA-1088.

**REQUEST A QUOTE**

Ready to request a quote for our Standardized Mite Extract? Or, have you ordered from us before? We now offer recurring orders that automatically ship your order in the quantity and frequency determined by your clinic's needs.

## SEE INSERT INSIDE

1. Arbes, Samuel J., Richard D. Cohn, Ming Yin, Michael L. Muilenburg, Harriet A. Burge, Warren Friedman and Darryl C. Zeldin. "House Dust Mite Allergen in U.S. Beds: Results from the First National Survey of Lead and Allergens in Housing." *Journal of Allergy and Clinical Immunology* 111.2 (2003): 408-14.

2. Jones, J. and Wallen, N. (1991). Comparison of puncture skin test (PST) reactivity to standardized D. farina (DF) extracts at 30,000 AU/ml (30k) and 10,000 AU/mL (10k). AAAAI.

## Why HollisterStier?

We believe in elevating allergy, together. Take advantage of our quality products, customized services, and personalized training now.



**HollisterStier**  
Allergy

Phone: 1.800.495.7437

Fax: 1.800.752.6258

Email: [HollisterStier@jubl.com](mailto:HollisterStier@jubl.com)

REQUEST A QUOTE



[HSALLERGY.COM](http://HSALLERGY.COM)